Temporal.pdf

This file contains additional information about the linear regression model that modeled weekly mosquito infection rate.

- A) Scatter plots between key covariates and mosquito infection
- B) Residual plots, autocorrelation function graphs, and Q-Q plots for the four linear regression models described in Table 2 in ms.
- C) Equations for each of the four models




Figure A. Scatterplots between mosquito infection rate (MIR), Degree Week (DW), precipitation with 3-week average (prcp3wk), and precipitation with 5-week average (prcp5wk).




Figure B1. Residuals from model 1 (top) and the Autocorrelation Function (ACF) for the residuals for each year (bottom)




Figure B2. Residuals from model 2 (top) and the Autocorrelation Function (ACF) for the residuals for each year (bottom)




Figure B3. Residuals from model 3 (top) and the Autocorrelation Function (ACF) for the residuals for each year (bottom)




Figure B4. Residuals from model 4 (top) and the Autocorrelation Function (ACF) for the residuals for each year (bottom)




Figure B5. Q-Q plot of the residuals in model 1 to 4 (from top to bottom). All years are combined in these plots.

Model 1

$$MIR(i,j) = 8.86 * DW(i-1,j) - 19.89 * prepann_abn(j-1) + 0.63624 * MIR(i-1,j)$$

Model 2

$$MIR(i,j) = 23.475 * DW(i-1,j) - 40.47 * prep annual(j-1) + 9.349 * prep3wk(i-3,j)$$

Model 3

$$MIR(i,j) = 11.4 * DW(i-4,j) + 0.6 * MIR(i-2,j) - 33.4 * prep_annual(j-1)$$

Model 4

$$MIR(i,j) = 32.5 * DW(i-4,j) + 11.4 * prep3wk(i-4,j) - 48.2 * prep_annual(j-1)$$

where the subscript i is the index for weeks, and j the index of year.

C) Equations for the four linear regression models described in the ms Table 2.